

A note on Closure Operators in Category of Modules

Vishvajit V S Gautam*

*The Institute of Mathematical Sciences
CIT Campus, Taramani, Chennai - 600113 INDIA*

Abstract

In this article we give application of closure operators in category of modules. Our main result shows that every subcategory $\mathcal{A}$ of injective modules of **R-mod** (under a mild condition) induces a torsion theory of **R-mod**.

AMS subject classification 2001 : 16D40

Keywords : closure operator, epimorphism, free ideal ring, monomorphism, regular closure operator, torsion theory.

Notion of closure operators (operations, systems, functions, relations) is known to us from algebra, logic, lattice theory and topology. Categorical view of closure operators play an important role in various branches of mathematics. In an arbitrary category $\mathcal{X}$ with suitable axiomatically defined notion of subobjects, a (categorical) closure operator c is defined to be a family $(c_X)_{X \in \mathcal{X}}$ satisfying the properties of extension, monotonicity and continuity. Concept of closure operators induced by a subcategory $\mathcal{A}$ of **Top** was introduced by Salbany in 1976 and have been studied by many researchers in a variety of situations. Closure operators are proved to be useful in study of Galois equivalence between certain factorization systems. In category of **R-mod** of R-modules closure operators correspond to preradicals. For more details, see [2].

In section 2 of this paper, we give characterizations of **R-mod** homomorphisms in terms of regular closure operators. Using these characterizations we establish that every subcategory $\mathcal{A}$ of injective modules of **R-mod** (under a mild condition) induces a torsion theory of **R-mod** [Theorem 2.3]. Some useful observations of these characterizations to modules over a *free ideal*

*vishvajit@imsc.res.in, gautamvvs@yahoo.com

rings(semifir or n-fir) are given in section 3.

Subcategories are always assumed to be full and isomorphism closed.

1 Preliminaries

Throughout this paper we consider a category $\mathcal{X}$ and a fixed class $\mathcal{M}$ of monomorphisms in $\mathcal{X}$ which contains all isomorphisms of $\mathcal{X}$. It is assumed that

- $\mathcal{M}$ is closed under composition;
- $\mathcal{X}$ is finite $\mathcal{M}$ -complete.

A *closure operator* c on the category $\mathcal{X}$ with respect to class $\mathcal{M}$ of subobjects is given by a family $c = (c_X)_{X \in \mathcal{X}}$ of maps $c_X : \mathcal{M}/X \rightarrow \mathcal{M}/X$ such that for every $X \in \mathcal{X}$

1. $m \leq c(m)$; 2. $m \leq m' \Rightarrow c(m) \leq c(m')$; and 3. for every $f : X \rightarrow Y$ and $m \in \mathcal{M}/X$, $f(c_X(m)) \leq c_Y(f(m))$.

For each $m \in \mathcal{M}$ we denote by $c(m)$ the c -closure of m .

An $\mathcal{M}$ -morphism $m \in \mathcal{M}/X$ is called c -closed if $m \cong c_X(m)$. A closure operator c is said to be idempotent if $c(c(m)) \cong c(m)$. In case $c(m \vee n) \cong c(m) \vee c(n)$ we say c to be additive. An $\mathcal{M}$ -subobject m of X is called c -dense in X if $c_X(m) \cong 1_X$.

For a subcategory $\mathcal{A}$ of $\mathcal{X}$, a morphism $f : X \rightarrow Y$ is an $\mathcal{A}$ -regular monomorphism if it is the equalizer of two morphisms $h, k : Y \rightarrow A$ with $A \in \mathcal{A}$.

Let $\mathcal{M}$ contain the class of regular monomorphisms of $\mathcal{X}$. For $m : M \rightarrow X$ in $\mathcal{M}$ define

$$c_{\mathcal{A}}(m) = \bigwedge \{r \in \mathcal{M} \mid r \geq m \text{ and } r \text{ is } \mathcal{A}\text{-regular}\}$$

which is a closure operator of $\mathcal{X}$. These closure operators are called regular and $c_{\mathcal{A}}(m)$ is called the $\mathcal{A}$ -closure of m . In case $\mathcal{A} = \mathcal{X}$ we denote $c_{\mathcal{A}}(m)$ by $c(m)$.

2 Closure Operators in category of modules

In this section we will see application of closure operators in category of modules.

Let $\mathcal{X} = \mathbf{R}\text{-mod}$ the category of R -modules and let $\mathcal{M}$ be the class of all monomorphisms of $\mathcal{X}$. In this Case $\mathcal{X}$ is $\mathcal{M}$ -complete.

Recall that by a linear function f on an R -module M we mean an R -module homomorphism $f : M \longrightarrow R$.

Let $\mathcal{A}$ be a subcategory of **R-mod**. For an object M of **R-mod**, $\mathcal{M}/M$ is the set of all submodules of M , we can identify each n in $\mathcal{M}/M$ with $N \subseteq M$ a submodule of M . Let N be a submodule of M . The $\mathcal{A}$ -regular closure operator of N can be computed by following formula ([2])

$$c_{\mathcal{A}}(N) = \cap \{Ker(g) \mid g \text{ is an } R\text{-module homomorphism from } M \text{ to } A \in \mathcal{A} \text{ and } g(N) = (0)\}.$$

Throughout the remainder of this paper $\mathcal{A}$ denotes a category of injective modules.

Proposition 2.1 $c_{\mathcal{A}}(N) = M$ i.e. N is $c_{\mathcal{A}}$ -dense in M if and only if $Hom_R(M/N, A) = 0$, $A \in \mathcal{A}$.

Proof. Let A be an object in $\mathcal{A}$. Consider the exact sequence

$$0 \longrightarrow N \longrightarrow M \longrightarrow M/N \longrightarrow 0$$

Since $Hom_R(-, A)$ is left exact functor, the sequence

$$0 \longrightarrow Hom_R(M/N, A) \longrightarrow Hom_R(M, A) \longrightarrow Hom_R(N, A) \longrightarrow 0$$

is exact. If $Hom_R(M/N, A) = 0$, above left exact sequence implies

$$Hom_R(N, A) = Hom_R(M, A).$$

Therefore every $f \in Hom_R(M, A)$ satisfying $f(N) = 0$ must be zero. Hence $c_{\mathcal{A}}(N) = Ker(0) = M$.

Conversely, assume that $f \in Hom_R(M/N, A)$ for some $A \in \mathcal{A}$. Let $\pi : M \longrightarrow M/N$ be the canonical projection. The composition $f \cdot \pi : M \longrightarrow A$ is an R -module homomorphism such that $f(\pi(n)) = f(N) = 0$ for all $n \in N$. This gives $c_{\mathcal{A}}(N) \subseteq Ker(f \cdot \pi)$ and $M = Ker(f \cdot \pi)$, which implies $f \cdot \pi = 0$, and so $f(\pi(M)) = 0(M) = 0$ implies $f(M/N) = 0$ and hence $f = 0$. $\square$

Proposition 2.2 $c_{\mathcal{A}}(N) = N$ i.e. N is $c_{\mathcal{A}}$ -closed in M if and only if $Hom_R(T/N, A) \neq 0$ for each non-zero submodule T/N of M/N and $A \in \mathcal{A}$.

Proof. Let $A \in \mathcal{A}$. If $c_{\mathcal{A}}(N) \neq N$, we can regard $c_{\mathcal{A}}(N)/N$ as a non-zero submodule of M/N . Suppose $0 \neq g \in Hom_R(c_{\mathcal{A}}(N)/N, A)$. Using the arguments similar to the previous Proposition we get $g(c_{\mathcal{A}}(N)/N) = 0$, which implies $g = 0$. A contradiction, therefore we must have $c_{\mathcal{A}}(N) = N$.

Conversely, if $c_{\mathcal{A}}(N) = N$, $c_{\mathcal{A}}(N)$ is not dense in M . Therefore by above proposition we have $\text{Hom}_R(M/N, A) \neq 0$. Let T/N be a non-zero submodule of M/N . Let $0 \neq f \in \text{Hom}_R(M/N, A)$. Then restriction of f to T/N gives $\text{Hom}_R(T/N, A) \neq 0$. $\square$

Next we relate the $\mathcal{A}$ -regular closure operator defined in the beginning of this section with torsion theory of **R-mod** in sense of [3].

Theorem 2.3 *Subcategory $\mathcal{A}$ of **R-mod** satisfying $\mathcal{T} \cap \mathcal{A} = 0$ for a class $\mathcal{T}$ of objects of **R-mod** induces a torsion theory for **R-mod** such that $\mathcal{A}$ forms a torsion free class consisting of torsion free objects.*

Proof. Consider the pair $(\mathcal{T}, \mathcal{F})$ of classes of subobject of **R-mod**, where

$$\mathcal{T} = \{M/N \mid c_{\mathcal{A}}(N) = M\}$$

and

$$\mathcal{F} = \mathcal{A}$$

It is immediate from above two propositions that $(\mathcal{T}, \mathcal{F})$ is a torsion theory. This torsion theory is also a *hereditary torsion theory*. $\square$

Let $(\mathcal{T}, \mathcal{F} = \mathcal{A})$ be a torsion theory induced by a subcategory $\mathcal{A}$ of **R-mod** in sense of Theorem 2.3 . Following result (cf. [3], Proposition 2.1, 2.2) characterizes the torsion classes.

Theorem 2.4 *For torsion theory $(\mathcal{T}, \mathcal{F} = \mathcal{A})$*

1. *The class $\mathcal{T}$ of objects of **R-mod** is closed under image, infinite sums and group extensions.*
2. *$\mathcal{A}$ is closed under kernel, infinite products and group extensions.*

3 Closure Operators in category of firs (semifirs or n -firs)

Recall that by a *free ideal ring* we mean a ring R with the property that all their (one-sided) ideal are free as left module (resp. right module) of unique rank.

Let **R-mod** denotes the category of modules whose underlying rings are *firs (semifirs or n -firs)* ([1]). Let R be injective as R -module. An R -module M is said to be bounded if $\text{Hom}_R(M, R) = 0$

Proposition 3.1 *Let N be a submodule of M .*

1. *A quotient module M/N is bounded if and only if $c_{\mathcal{A}}(N) = M$.*
2. *A quotient module M/N is unbounded if and only if $c_{\mathcal{A}}(N) = N$.*

Proof. Obvious. $\square$

Clearly bounded injective modules form a torsion class in above sense.

Proposition 3.2 1. *$c_{\mathcal{A}}(N) = N$ if and only if M/N has a direct summand which is free of positive rank.*

2. *$c_{\mathcal{A}}(N) = M$ if and only if M/N does not have a direct summand which is free of positive rank.*

Proof. 1. Since the image of every non-zero homomorphism f is free, which implies M/N has a direct summand.

Conversely, If M/N is a direct summand of a free module $f(M/N)$ of positive rank. Then f should be free.

2. The result is immediate from 1. $\square$

Many fragment of torsion theory for **R-mod** can be related with results proved in this section. Theorem 2.3 provides a convenient way to do this. One can use these result in the study of Gabriel topology [3]. If we assume R to be *fir*(*semifir*, *n-fir* or Bezout domain), there are some areas of *free ideal ring theory* where these results are useful. It may be a subject of further investigation.

References

- [1] P.M. Cohn, *Free rings and their relations* (Academic Press, London, 1971).
- [2] D. Dikranjan and W. Tholen, *Categorical structure of closure operators with applications to Topology, Algebra and Discrete Mathematics* (Kluwer, Dordrecht, 1994).
- [3] B. Stenström, *Rings of quotients* (Springer-Verlag, Berlin, 1975)